

## Homework 4

Due: December 6, 2015

**Note:** You may discuss these problems in groups. However, you must write up your own solutions and mention the names of the people in your group. Also, please do mention any books, papers or other sources you refer to. It is recommended that you typeset your solutions in L<sup>A</sup>T<sub>E</sub>X.

## 1. Gaussian Random Variables.

[5+5+5]

Prove the following very useful facts about Gaussian random variables:

- (a) Let  $\mathbf{u}, \mathbf{v} \in \mathbb{R}^n$  be two vectors. Let  $\mathbf{g} \in \mathbb{R}^n$  be a random vector such that each coordinate  $g_i$  of  $\mathbf{g}$  is distributed as a Gaussian random variable with mean 0 and variance 1, and any two coordinates  $g_i, g_j$  (for  $i \neq j$ ) are independent. Then show that

$$\mathbb{E}_{\mathbf{g}}[\langle \mathbf{u}, \mathbf{g} \rangle \cdot \langle \mathbf{v}, \mathbf{g} \rangle] = \langle \mathbf{u}, \mathbf{v} \rangle .$$

- (b) Let  $g$  be a Gaussian random variable with mean 0 and variance 1. Show that for any  $t \in \mathbb{R}$ , we have

$$\mathbb{E}[e^{tg}] = e^{t^2/2} .$$

Comparing coefficients of  $t^{2k}$  on both sides, use this to show that for any  $k \in \mathbb{N}$ ,

$$\mathbb{E}[g^{2k}] = \frac{(2k)!}{2^k \cdot k!} .$$

- (c) Let  $g_1, g_2, g_3$  and  $g_4$  be (not necessarily independent) Gaussian random variables with mean 0. Additionally, assume that for *all* coefficients  $\alpha_1, \dots, \alpha_4 \in \mathbb{R}$ , the linear combination  $\alpha_1 g_1 + \dots + \alpha_4 g_4$  is also a Gaussian random variable (note that this is not always true if  $g_1, \dots, g_4$  are not independent). Consider the function  $\mathbb{E}_{g_1, g_2, g_3, g_4}[e^{t_1 g_1 + t_2 g_2 + t_3 g_3 + t_4 g_4}]$  in the variables  $t_1, t_2, t_3, t_4$  and use it to show that

$$\mathbb{E}[g_1 g_2 g_3 g_4] = \mathbb{E}[g_1 g_2] \cdot \mathbb{E}[g_3 g_4] + \mathbb{E}[g_1 g_3] \cdot \mathbb{E}[g_2 g_4] + \mathbb{E}[g_1 g_4] \cdot \mathbb{E}[g_2 g_3] .$$

This shows that for *any* four Gaussian random variables, the expectation of their product can be expressed in terms of their pairwise correlations! This is a special case of what is known as Wick's theorem, which can also be proved by the above method.

## 2. Spectra of Bipartite Graphs.

[2+3+5]

Let  $G = (U, V, E)$  be a  $d$ -regular bipartite graph with adjacency matrix  $A$ , where  $U, V$  represent the two sides of the graph.

- (a) Prove that for the graph to be regular, we must have  $|U| = |V|$ .  
 (b) Prove that  $-d$  is an eigenvalue of  $A$  and find the corresponding eigenvector.  
 (c) Prove that for every eigenvalue  $\mu$  of  $A$ ,  $-\mu$  is also an eigenvalue of  $A$ .