

Lecture 1: October 1, 2019

Lecturer: Madhur Tulsiani

The primary goal of this course is to collect a set of basic mathematical tools which are often useful in various areas of computer science. We will mostly focus on various applications of linear algebra and probability. Please see the course webpage for a more detailed list of topics.

The course will be evaluated on the basis of the following:

- Homeworks: 40% (four homeworks contributing 10% each)
- Quizzes: 10% (two quizzes contributing 5% each)
- Midterm: 20%
- Final: 30%

We will spend 3-4 of lectures reviewing some of the basic concepts of linear algebra before we move on to some of the applications.

Here's a couple of problems to think about if you are already familiar with the contents of this lecture. The first is taken Laci Babai's REU course and the second from the excellent book "Thirty Three Miniatures" by Jiří Matoušek [Mat10], both which I highly recommend for many more fun applications of Linear Algebra. Both the materials are linked from the "resources" section of the course webpage.

Problem 0.1 Let w_1, \dots, w_{13} be thirteen positive real weights such that when any one weight is removed, the other 12 can be partitioned into two sets of six weights each, such that the total weights of the two sets are equal. Prove that all thirteen of the weights must be equal.

Problem 0.2 Consider the problem of covering the binary hypercube $\{0, 1\}^n$ using hyperplanes, which are subsets of \mathbb{R}^n of the form

$$\mathcal{H} := \left\{ x \in \mathbb{R}^n \mid \sum_{i=1}^n a_i \cdot x_i = b \right\},$$

where $a_1, \dots, a_n, b \in \mathbb{R}$. Note that the two hyperplanes $x_1 = 0$ and $x_1 = 1$ suffice to cover the entire set $\{0, 1\}^n$. However, suppose we want to cover the set $\{0, 1\}^n \setminus \{0^n\}$ only using hyperplanes not containing the origin. Show that the number of hyperplanes needed is at least n . Is this tight?

1 Fields

A field, often denoted by \mathbb{F} , is simply a nonempty set with two associated operations $+$ and \cdot mapping $\mathbb{F} \times \mathbb{F} \rightarrow \mathbb{F}$, which satisfy:

- **commutativity:** $a + b = b + a$ and $a \cdot b = b \cdot a$ for all $a, b \in \mathbb{F}$.
- **associativity:** $a + (b + c) = (a + b) + c$ and $a \cdot (b \cdot c) = (a \cdot b) \cdot c$ for all $a, b, c \in \mathbb{F}$.
- **identity:** There exist elements $0_{\mathbb{F}}, 1_{\mathbb{F}} \in \mathbb{F}$ such that $a + 0_{\mathbb{F}} = a$ and $a \cdot 1_{\mathbb{F}} = a$ for all $a \in \mathbb{F}$.
- **inverse:** For every $a \in \mathbb{F}$, there exists an element $(-a) \in \mathbb{F}$ such that $a + (-a) = 0_{\mathbb{F}}$. For every $a \in \mathbb{F} \setminus \{0_{\mathbb{F}}\}$, there exists $a^{-1} \in \mathbb{F}$ such that $a \cdot a^{-1} = 1_{\mathbb{F}}$.
- **distributivity of multiplication over addition:** $a \cdot (b + c) = a \cdot b + a \cdot c$ for all $a, b, c \in \mathbb{F}$.

Example 1.1 \mathbb{Q} , \mathbb{R} and \mathbb{C} with the usual definitions of addition and multiplication over these fields.

Example 1.2 Consider defining addition and multiplication on \mathbb{Q}^2 as

$$(a, b) + (c, d) = (a + c, b + d) \quad \text{and} \quad (a, b) \cdot (c, d) = (ac + bd, ad + bc).$$

These operations do not define a field. While various properties of addition are indeed satisfied, inverses may not always exist for multiplication as defined above. Check that the multiplicative identity needs to be $(1, 0)$ but then the element $(1, -1)$ has no multiplicative inverse.

However, for any prime p , the following operations do define a field

$$(a, b) + (c, d) = (a + c, b + d) \quad \text{and} \quad (a, b) \cdot (c, d) = (ac + pbd, ad + bc).$$

This is equivalent to taking $\mathbb{F} = \{a + b\sqrt{p} \mid a, b \in \mathbb{Q}\}$ with the same notion of addition and multiplication as for real numbers. Alternatively, one can also define a field by taking $(a, b) \cdot (c, d) = (ac - bd, ad + bc)$ (why?)

Example 1.3 For any prime p , the set $\mathbb{F}_p = \{0, 1, \dots, p - 1\}$ (also denoted as $GF(p)$) is a field with addition and multiplication defined modulo p .

Exercise 1.4 Show that the set $\{a + b\sqrt[3]{2} + c\sqrt[3]{4} \mid a, b, c \in \mathbb{Q}\}$ is a field.

2 Vector Spaces

A vector space V over a field \mathbb{F} is a nonempty set with two associated operations $+$: $V \times V \rightarrow V$ (vector addition) and \cdot : $\mathbb{F} \times V \rightarrow V$ (scalar multiplication) which satisfy:

- **commutativity of addition:** $v + w = w + v$ for all $v, w \in V$.
- **associativity of addition:** $u + (v + w) = (u + v) + w \forall u, v, w \in V$.
- **pseudo-associativity of scalar multiplication:** $a \cdot (b \cdot v) = (a \cdot b) \cdot v \forall a, b \in \mathbb{F}, v \in V$.
- **identity for vector addition:** There exists $0_V \in V$ such that for all $v \in V$, $v + 0_V = v$.
- **inverse for vector addition:** $\forall v \in V, \exists (-v) \in V$ such that $v + (-v) = 0_V$.
- **distributivity:** $a \cdot (v + w) = a \cdot v + a \cdot w$ and $(a + b) \cdot v = a \cdot v + b \cdot v$ for all $a, b \in \mathbb{F}$ and $v, w \in V$.
- **identity for scalar multiplication:** $1_{\mathbb{F}} \cdot v = v$ for all $v \in V$.

Example 2.1 Any field \mathbb{F} is a vector space over itself.

Example 2.2 \mathbb{R} is a vector space over \mathbb{Q} .

Example 2.3 $\mathbb{R}[X]$ is a vector space over \mathbb{R} .

Example 2.4 $C([0, 1], \mathbb{R}) = \{f : [0, 1] \rightarrow \mathbb{R} \mid f \text{ is continuous}\}$ is a vector space over \mathbb{R} .

Example 2.5 $\text{Fib} = \{f \in \mathbb{R}^{\mathbb{N}} \mid f(n) = f(n-1) + f(n-2) \forall n \geq 2\}$ is a vector space over \mathbb{R} .

Definition 2.6 (Linear Dependence) A set $S \subseteq V$ is linearly dependent if there exist distinct $v_1, \dots, v_n \in S$ and $a_1, \dots, a_n \in \mathbb{F}$ not all zero, such that $\sum_{i=1}^n a_i \cdot v_i = 0_V$. A set which is not linearly dependent is said to be linearly independent.

Example 2.7 The set $\{1, \sqrt{2}, \sqrt{3}\}$ is linearly independent in the vector space \mathbb{R} over the field \mathbb{Q} .

Exercise 2.8 Let $a_1, \dots, a_n \in \mathbb{R}$ be distinct and let $g(x) = \prod_{i=1}^n (x - a_i)$. Define

$$f_i(x) = \frac{g(x)}{x - a_i} = \prod_{j \neq i} (x - a_j),$$

where we extend the function at point a_i by continuity. Prove that f_1, \dots, f_n are linearly independent in the vector space $\mathbb{R}[x]$ over the field \mathbb{R} .

Exercise 2.9 *Prove that the set of functions*

$$S = \{1\} \cup \{\sin(kx) \mid k \in \mathbb{N}, k \geq 1\} \cup \{\cos(kx) \mid k \in \mathbb{N}, k \geq 1\},$$

is linearly independent in the vector space of continuous real-valued functions over \mathbb{R} .

References

[Mat10] Jiří Matoušek, *Thirty-three miniatures: Mathematical and algorithmic applications of linear algebra*, vol. 53, American Mathematical Soc., 2010. [1](#)