

Lecture 1: September 26, 2023

Lecturer: Madhur Tulsiani

The primary goal of this course is to collect a set of basic mathematical tools which are often useful in various areas of computer science. We will mostly focus on various applications of linear algebra and probability. Please see the course webpage for a more detailed list of topics.

The course will be evaluated on the basis of the following:

- Homeworks: 60% (five homeworks contributing 12% each)
- Midterm: 15%
- Final: 25%

We will spend 3-4 of lectures reviewing some of the basic concepts of linear algebra before we move on to some of the applications.

Here's a couple of problems to think about if you are already familiar with the contents of this lecture. These are from the excellent book "Thirty Three Miniatures" by Jiří Matoušek [Mat10], both which I highly recommend for many more fun applications of Linear Algebra (you can find a link in the "resources" section of the course webpage.)

Problem 0.1 *Show that a rectangle with sides 1 and $\sqrt{2}$ cannot be tiled with a finite number of non-overlapping squares. In fact, you can try to prove that this is the case when $\sqrt{2}$ is replaced by any irrational number x .*

Problem 0.2 *Can there be four points in the plane such that all pairs of distances are odd integers?*

1 Fields

A field, often denoted by \mathbb{F} , is simply a nonempty set with two associated operations $+$ and \cdot mapping $\mathbb{F} \times \mathbb{F} \rightarrow \mathbb{F}$, which satisfy:

- **commutativity:** $a + b = b + a$ and $a \cdot b = b \cdot a$ for all $a, b \in \mathbb{F}$.
- **associativity:** $a + (b + c) = (a + b) + c$ and $a \cdot (b \cdot c) = (a \cdot b) \cdot c$ for all $a, b, c \in \mathbb{F}$.

- **identity:** There exist elements $0_{\mathbb{F}}, 1_{\mathbb{F}} \in \mathbb{F}$ such that $a + 0_{\mathbb{F}} = a$ and $a \cdot 1_{\mathbb{F}} = a$ for all $a \in \mathbb{F}$.
- **inverse:** For every $a \in \mathbb{F}$, there exists an element $(-a) \in \mathbb{F}$ such that $a + (-a) = 0_{\mathbb{F}}$. For every $a \in \mathbb{F} \setminus \{0_{\mathbb{F}}\}$, there exists $a^{-1} \in \mathbb{F}$ such that $a \cdot a^{-1} = 1_{\mathbb{F}}$.
- **distributivity of multiplication over addition:** $a \cdot (b + c) = a \cdot b + a \cdot c$ for all $a, b, c \in \mathbb{F}$.

Example 1.1 \mathbb{Q}, \mathbb{R} and \mathbb{C} with the usual definitions of addition and multiplication over these fields.

Example 1.2 Consider defining addition and multiplication on \mathbb{Q}^2 as

$$(a, b) + (c, d) = (a + c, b + d) \quad \text{and} \quad (a, b) \cdot (c, d) = (ac + bd, ad + bc).$$

These operations do not define a field. While various properties of addition are indeed satisfied, inverses may not always exist for multiplication as defined above. Check that the multiplicative identity needs to be $(1, 0)$ but then the element $(1, -1)$ has no multiplicative inverse.

However, for any prime p , the following operations do define a field

$$(a, b) + (c, d) = (a + c, b + d) \quad \text{and} \quad (a, b) \cdot (c, d) = (ac + pbd, ad + bc).$$

This is equivalent to taking $\mathbb{F} = \{a + b\sqrt{p} \mid a, b \in \mathbb{Q}\}$ with the same notion of addition and multiplication as for real numbers. Alternatively, one can also define a field by taking $(a, b) \cdot (c, d) = (ac - bd, ad + bc)$ (why?)

Example 1.3 For any prime p , the set $\mathbb{F}_p = \{0, 1, \dots, p - 1\}$ (also denoted as $GF(p)$) is a field with addition and multiplication defined modulo p . Also, check that defining addition and multiplication modulo a composite number (say modulo 6) does not give a field.

Exercise 1.4 Show that the set $\{a + b\sqrt[3]{2} + c\sqrt[3]{4} \mid a, b, c \in \mathbb{Q}\}$ is a field.

2 Vector Spaces

A vector space V over a field \mathbb{F} is a nonempty set with two associated operations $+$: $V \times V \rightarrow V$ (vector addition) and \cdot : $\mathbb{F} \times V \rightarrow V$ (scalar multiplication) which satisfy:

- **commutativity of addition:** $v + w = w + v$ for all $v, w \in V$.
- **associativity of addition:** $u + (v + w) = (u + v) + w \forall u, v, w \in V$.

- **pseudo-associativity of scalar multiplication:** $a \cdot (b \cdot v) = (a \cdot b) \cdot v \forall a, b \in \mathbb{F}, v \in V$.
- **identity for vector addition:** There exists $0_V \in V$ such that for all $v \in V, v + 0_V = v$.
- **inverse for vector addition:** $\forall v \in V, \exists (-v) \in V$ such that $v + (-v) = 0_V$.
- **distributivity:** $a \cdot (v + w) = a \cdot v + a \cdot w$ and $(a + b) \cdot v = a \cdot v + b \cdot v$ for all $a, b \in \mathbb{F}$ and $v, w \in V$.
- **identity for scalar multiplication:** $1_{\mathbb{F}} \cdot v = v$ for all $v \in V$.

Example 2.1 Any field \mathbb{F} is a vector space over itself.

Example 2.2 \mathbb{R} is a vector space over \mathbb{Q} .

Example 2.3 Let $\mathbb{R}[x]$ denote the space of polynomials in one variable x , with real coefficients i.e.,

$$\mathbb{R}[x] := \left\{ \sum_{i=0}^t c_i \cdot x^i \mid t \in \mathbb{N}, c_0, \dots, c_t \in \mathbb{R} \right\}.$$

Then, $\mathbb{R}[x]$ is a vector space over \mathbb{R} . Similarly, the space $\mathbb{R}^{\leq d}[x]$ of polynomials with degree at most d , defined as

$$\mathbb{R}^{\leq d}[x] := \left\{ \sum_{i=0}^t c_i \cdot x^i \mid t \in \mathbb{N}, t \leq d, c_0, \dots, c_t \in \mathbb{R} \right\},$$

is also a vector space over \mathbb{R} .

Example 2.4 $C([0, 1], \mathbb{R}) = \{f : [0, 1] \rightarrow \mathbb{R} \mid f \text{ is continuous}\}$ is a vector space over \mathbb{R} .

Example 2.5 $\text{Fib} = \{f \in \mathbb{R}^{\mathbb{N}} \mid f(n) = f(n-1) + f(n-2) \forall n \geq 2\}$ is a vector space over \mathbb{R} .

Definition 2.6 (Linear Dependence) A set $S \subseteq V$ is linearly dependent if there exist distinct $v_1, \dots, v_n \in S$ and $c_1, \dots, c_n \in \mathbb{F}$ not all zero, such that $\sum_{i=1}^n c_i \cdot v_i = 0_V$. A set which is not linearly dependent is said to be linearly independent. A sum of the form $\sum_{i=1}^n c_i \cdot v_i$ is referred to as a linear combination of the vectors v_1, \dots, v_n .

Example 2.7 The set $\{1, \sqrt{2}, \sqrt{3}\}$ is linearly independent in the vector space \mathbb{R} over the field \mathbb{Q} .

Exercise 2.8 Let $a_1, \dots, a_n \in \mathbb{R}$ be distinct and let $g(x) = \prod_{i=1}^n (x - a_i)$. Define

$$f_i(x) = \frac{g(x)}{x - a_i} = \prod_{j \neq i} (x - a_j),$$

where we extend the function at point a_i by continuity. Prove that f_1, \dots, f_n are linearly independent in the vector space $\mathbb{R}[x]$ over the field \mathbb{R} .

3 Span and Bases

Definition 3.1 Given a set $S \subseteq V$, we define its span as

$$\text{Span}(S) = \left\{ \sum_{i=1}^n a_i \cdot v_i \mid a_1, \dots, a_n \in \mathbb{F}, v_1, \dots, v_n \in S, n \in \mathbb{N} \right\}.$$

Note that we only include finite linear combinations. Also, since linear combinations of vectors are still in V , we have $\text{Span}(S) \subseteq V$. In fact, you can check that $\text{Span}(S)$ is also a vector space. Such a subset of V , which is also a vector space, is called a subspace of V .

Remark 3.2 Note that the definition above and the previous definitions of linear dependence and independence, all involve only finite linear combinations of the elements. Infinite sums cannot be said to be equal to a given element of the vector space without a notion of convergence or distance, which is not necessarily present in an abstract vector space.

Definition 3.3 A set B is said to be a basis for the vector space V if B is linearly independent and $\text{Span}(B) = V$.

3.1 Lagrange interpolation

Lagrange interpolation is used to find the unique polynomial of degree at most $n - 1$, taking given values at n distinct points. We can derive the formula for such a polynomial using basic linear algebra.

Let $a_1, \dots, a_n \in \mathbb{R}$ be distinct. Say we want to find the unique (why?) polynomial p of degree at most $n - 1$ satisfying $p(a_i) = b_i \forall i \in [n]$. Recall that the space of polynomials of degree at most $n - 1$ with real coefficients, denoted by $\mathbb{R}^{\leq n-1}[x]$, is a vector space. Also, recall from the last lecture that if we define $g(x)$ as $\prod_{i=1}^n (x - a_i)$, the degree $n - 1$ polynomials defined as

$$f_i(x) = \frac{g(x)}{x - a_i} = \prod_{j \neq i} (x - a_j),$$

are n linearly independent polynomials in $\mathbb{R}^{\leq n-1}[x]$. Thus, they must form a basis for $\mathbb{R}^{\leq n-1}[x]$ and we can write the required polynomial, say p as

$$p = \sum_{i=1}^n c_i \cdot f_i,$$

for some $c_1, \dots, c_n \in \mathbb{R}$. Evaluating both sides at a_i gives $p(a_i) = b_i = c_i \cdot f_i(a_i)$. Thus, we get

$$p(x) = \sum_{i=1}^n \frac{b_i}{f_i(a_i)} \cdot f_i(x).$$

Exercise 3.4 Check that the above argument can be used to find a polynomial of degree at most $n - 1$ in the space $\mathbb{F}[x]$ for any field \mathbb{F} such that $|\mathbb{F}| \geq n$.

3.2 Secret Sharing

Consider the problem of sharing a secret s , which is an integer in a known range $[0, M]$ with a group of n people, such that if any d of them get together, they are able to learn the secret message. However, if fewer than d of them are together, they do not get any information about the secret. We can then proceed as follows:

- Choose a finite field \mathbb{F}_p , with $p > \max(n, M)$.
- Choose $d - 1$ random values b_1, \dots, b_{d-1} in $\{0, \dots, p - 1\}$, and let $Q \in \mathbb{F}_p^{\leq d-1}[x]$ be the polynomial

$$Q = s + b_1x + b_2x^2 + \dots + b_{d-1}x^{d-1}.$$

Note that the secret is $Q(0)$.

- For $i = 1, \dots, n$, give person i the pair $(i, Q(i))$.

Note that if any group of d or more people get together, they can uniquely determine the polynomial Q by Lagrange interpolation. They can then recover the secret by evaluating Q at 0. However, if $d - 1$ of them gather, then there is always a polynomial consistent with the values they hold, and any possible value at 0. To precisely say that they learn nothing about the secret, we use the fact that there is *exactly one* polynomial consistent with the values they hold and any given value at 0. Since for any given secret s there are exactly p^{d-1} polynomials with $Q(0) = s$, and we chose the polynomial at random conditioned on the secret, this means that any two secrets have the same probability of producing the observed $(d - 1)$ -tuple of shares. We will talk in more depth about arguments like this when we discuss probability in the second half of the course.

References

[Mat10] Jiří Matoušek, *Thirty-three miniatures: Mathematical and algorithmic applications of linear algebra*, vol. 53, American Mathematical Soc., 2010. 1